



ORAL PRESENTATION

Open Access

HTLV-3 and HTLV-4 antisense proteins enhance the transactivation potential of several Jun family members through interaction via their bZIP-like domain

Émilie Larocque^{1*}, Charlotte André-Arpin², Jean-Michel Mesnard², William M Switzer³, Benoit Barbeau¹

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses
Montreal, Canada. 26-30 June 2013

HTLV-3 and HTLV-4 are two recently identified viruses closely related to HTLV-1, which have not yet been linked to any diseases. Similarly to HTLV-1 and HTLV-2, which both express antisense transcript-derived HBZ and APH-2, respectively, HTLV-3 and HTLV-4 harbor antisense genes termed APH-3 and APH-4. These two encoded proteins contain an atypical bZIP domain but are still able to down-regulate Tax-mediated LTR activation, alike HBZ and APH-2. Since HBZ and APH-2 affect Jun-dependent transcription differently, our goal was to assess the impact of APH-3 and APH-4 on the transactivation potential of Jun family members. Co-IP experiments first showed that both APH-3 and APH-4 interacted with all tested Jun members. Jun/APH complexes significantly enhanced Jun-dependent transactivation of the human collagenase promoter containing a single AP-1 binding site as well as a construct bearing a minimal promoter and AP-1-binding sites. APH-3 and APH-4 deletion mutants and point mutations of specific leucine residues demonstrated that this functional interaction was mediated by the bZIP-like domain. Using a construct with a minimal promoter bearing GAL4-binding sites and APH-3/APH-4 expression vectors fused to the GAL4 DNA binding domain, we further showed that, in contrast to HBZ, APH-3 and APH-4 did not contain an activation domain. These results highlight the varying capacity of different HTLV-encoded antisense proteins to act upon transactivation mediated by Jun transcription factors. Taking into account the

previously reported association between HBZ and ATL development, determining the functional differences of these HTLV antisense proteins could contribute in better understanding this association.

Authors' details

¹Département des Sciences Biologiques and Centre de Recherche BioMed, Université du Québec à Montréal, Montréal (Québec) Canada. ²Université Montpellier 1, Centre d'Etudes d'Agents Pathogènes et Biotechnologies pour la Santé, CNRS, UM5236, Montpellier, France. ³Laboratory Branch, Division of HIV/AIDS Prevention, National Center for HIV, STD, and TB prevention, Centers for Disease Control and Prevention, Atlanta, GA, USA.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-O72

Cite this article as: Larocque et al.: HTLV-3 and HTLV-4 antisense proteins enhance the transactivation potential of several Jun family members through interaction via their bZIP-like domain. *Retrovirology* 2014 **11**(Suppl 1):O72.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: larocque.emilie@gmail.com

¹Département des Sciences Biologiques and Centre de Recherche BioMed, Université du Québec à Montréal, Montréal (Québec) Canada
Full list of author information is available at the end of the article

© 2014 Larocque et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.